

# A COMPARATIVE STUDY OF THE FLOWERING PHENOLOGY OF WILD GROWING GEOPHYTES IN TWO DIFFERENT MESOCLIMATIC AREAS IN THE CARPATHIAN BASIN

Krisztina Verbényiné Neumann<sup>1\*</sup>, Tivadar Baltazár<sup>2</sup>, Sarolta Meinhardt<sup>1</sup>,  
Orsolya Szirmai<sup>3</sup>

<sup>1\*</sup>Doctoral School of Environmental Sciences, Hungarian University of Agriculture and Life Sciences. H-2100 Páter Károly street 1, Gödöllő, HUNGARY, neumann.krisztina86@gmail.com

<sup>2</sup>Department of Agrochemistry, Soil Science, Microbiology and Plant Nutrition, Faculty of AgriSciences, Mendel University in Brno, Zemědělská 1665/1, 613 00 Brno, Czech Republic; baltazartivadar@gmail.com

<sup>3</sup>Institute of Animal Sciences and Wildlife Management, Faculty of Agriculture, University of Szeged, Andrassy út 15, H-6800 Hódmezővásárhely, Hungary, szirmai.orsolya@szte.hu

\*corresponding author: neumann.krisztina86@gmail.com

**Abstract:** Plant phenology - timing of cyclical or seasonal biological events - has proven to be a very sensitive indicator for climate change impacts. The phenology of many plant species has been advanced by warming, with earlier spring species being more sensitive. To understand better the driving factors of the changing phenology we investigated the phenology of different wild growing geophytes in the Carpathian Basin for three consecutive years. The study has been carried out as an *ex situ* experiment in two different mesoclimatic sites, one in the Gödöllő Botanical Garden of the Hungarian University of Agriculture and Life Sciences (average temperature 11,35 °C), other in the Eötvös Loránd University Botanical Garden, Fűvészkert in the central part of Budapest (average temperature 13,16 °C). During the experiment 5 replicates of 5 wild growing geophytes were examined. The results show an advance of 3.64 days in Budapest, with strong variation across species. The earliest flowering species didn't bloom (*Galanthus nivalis*) or died (*Eranthis hyemalis*) by the 3<sup>rd</sup> year in the site Budapest. If global warming continues, this advance and negative effects on wild growing plant species might be more serious in the future.

**Keywords:** Reproductive phenology, ex situ, climate change, heat island, botanical garden, temperate zone, early-spring.

## 1. Introduction

Climate change rises global temperatures, thus influencing ecosystem processes (Peñuelas 2017). On average, the world has already warmed 1.1 °C, affecting natural ecosystems in Europe and everywhere on Earth (IPCC, 2022). The observed trend of warming at a global or local scale can have serious implications on living organisms. Warming will decrease suitable habitat area for current terrestrial ecosystems and change their composition (IPCC, 2022). In Europe, more than half of the vascular plant flora may become endangered by the year 2080 as a result of

climatic changes (Thuiller et al. 2005). Based on current research it seems that climate change can no longer be stopped. Therefore, it is crucial to investigate possible adaptations (Li et al. 2019).

Plant phenology, the timing of seasonally recurring phenomena in plants (Schwartz 2013) has proven to be a very sensitive indicator for climate change impacts (Sparks 2002, Cleland 2007). Climate can strongly influence phenology by speeding up or delaying events such as emergence, peak activity and reproduction (Sherry et al. 2007, Wolkovich 2021). Phenology of many plant species has been already advanced by warming (Sparks et al. 2000, Fitter and Fitter 2002, Parmesan and Yohe 2003, Elzinga et al. 2007, Bertin 2008, Szabó et al. 2016, Neumann & Czóbel 2021). Changes in the plant reproductive period also have important consequences on the reproductive success of populations, and thus on their dynamics (Sherry et al. 2007). For example, changes in flowering time may disrupt plant–pollinator interactions, particularly when the pollinators are seasonal (e.g. insects), and reduce seed production of plants and food resources to the pollinators, thereby influencing the survival and success of both species (Fitter & Fitter, 2002). Many studies have reported species-specific phenological shifts in response to climate change (Forrest & Miller- Rushing, 2010; Renner & Zohner, 2018) and found that phenological timing in early-spring bloomers were more responsive to warming than mid- or late-spring bloomers (Fitter and Fitter 2002, Menzel et al., 2006; Miller-Rushing & Primack, 2008, Kubov et al 2022). Insect-pollinated plants tended to have greater advancement of flowering than wind-pollinated plants (Fitter and Fitter, 2002), annuals demonstrated greater advancement than perennials (Fitter and Fitter, 2002).

Many studies in middle and high latitudes demonstrate that the temperature is the main driving force and interannual modulator of phenological change, while other factors (e.g. photoperiod, precipitation etc.) only play a secondary role as limiting factors (Szabó et al. 2016, Wolkovich 2021). Among several environmental factors (light, moisture, temperature) that can affect bulb development, temperature has been established as playing a predominant role in controlling growth and flowering in bulbs (Khadorova and Boitel-Conti 2013). Besides that, for early spring geophytes the number of frosty days and snowmelt has an important effect on both leafing and flowering phenology, controlling spring soil temperature and moisture (Eppich et al 2009, Snopková & Hýrošová 2017, Bandoc et al. 2022).

Temperature sensitivity or phenological sensitivity, which is expressed as the date of phenological event change for per degree Celsius change of temperature (days /°C), has been widely used to characterize the plants' responses to changed temperature (Wang et. al 2015, Zhang et al. 2015). Since the temperature sensitivity of plant phenological stages determines the magnitude of phenological shifts in response to future climate warming, more attention has been paid to it, both in observational records and warming experiment studies (Wolkovich et al. 2012).

Slow-colonizing forest understorey plants are probably not able to rapidly adjust their distribution range following largescale climate change. Therefore, the acclimation potential to climate change within their actual occupied habitats will

likely be key for their short- and long-term persistence (deFrenne et al. 2010). This makes it particularly important to study the responses of these species to climate change. Nevertheless, relatively few studies have addressed the phenological events of geophytes as a life form (Tarasjev 1997, Turisova et al. 2007, Eppich et al 2009, Thomson 2010, Khodorova and Boitel-Conti 2013, Szabó et al. 2016, Snopková & Hýrošová 2017, Crişan et al. 2018, Puchalka et al. 2022).

Shifts in the phenological events of geophytes in the Carpathian Basin (Central Europe) are particularly poorly documented, with a few exceptions coming from the works of Eppich 2009, Szabó et al 2016, Neumann & Czóbel 2021).

Urban climate conditions are considered similar to the changing global climate conditions; therefore, many researchers study urbanized areas as small-scale experiments, or models, of global climate change (Ziska et al., 2003). Thus the urban environment is suitable for the application of the Space for Time substitution method. Which method encompasses analyses in which contemporary spatial phenomena are used to understand and model temporal processes that are otherwise unobservable, most notably past and future events. This method is used to predict the effects of climate change on biodiversity, identifying general trends, therefore its application saves time and money compared to long-term studies. (Pickett 1989, Blous et al. 2013) Thus, it is key to examine the patterns and shifts in the patterns of flowering phenology in urban areas compared with rural ones. Cities are strongly affected by climate change.

As part of the research, we examined the phenology of 5 geophyte species wild growing in the Carpathian Basin in two areas with different mesoclimates, in an ex situ experiment. We assumed (i) that first flowering, and the end of flowering will occur earlier at the site with a higher average temperature, while (ii) the length of flowering will longer the in the area with a higher average temperature. Furthermore, we examined to what temperature and other climatic factors besides the average annual temperature influence the time of occurrence of the different phenophases.

## **2. Materials and methods**

### **2.1. Study sites**

The study was carried out in two different mesoclimatic sites, one of which located in the Gödöllő Botanical Garden of Hungarian University of Agriculture and Life Sciences (47°35'36.2"N 19°22'06.2"E, 250 m elevation) (Szirmai et al. 2014), while the other in the Eötvös Loránd University Botanical Garden in the central part of Budapest (Budapest 47°29'05.6"N 19°05'05.7"E, 114 m elevation) (Orlóci et al 2019). Within a radius of 250 m around the two botanical gardens, the following local climate zones (LCZ) are present. In Budapest: LCZ 5 - Open mid-rise 60%, LCZ 6 - Open low rise 20% LCZ 2 - Compact mid-rise 20%. In Gödöllő: LCZ A - dense trees 40%, LCZ D - low plants 50%, LCZ 6 - Open low-rise 10% (Stewart & Oke 2012, [http1](http://)). The Gödöllő site is next to a natural forest fragment, part of the 4.5 hectar botanical garden. The Budapest site is located in the 3,1 hectar Fűvészkert.

During the three-year experiment the average air temperature was 11.35 °C and the average annual precipitation was 475.1 mm in Gödöllő Botanical Garden, while

in the Eötvös Loránd University Botanical Garden the average air temperature was 13.16 °C and an average annual precipitation was 527.4 mm. There was therefore a difference of 1.81 °C between the three-year average temperature of the two botanical gardens. We used this value to calculate the phenological sensitivity. A homogeneous patch was created for each selected species in the two sites.

## 2.2. Methods

To examine the effect of the different mesoclimatic environments on the phenology of geophyte species. We selected 5 different, (in the Carpathian Basin wild growing geophyte species: two early-spring bloomers *Galanthus nivalis* L., *Eranthis hyemalis* L., and three mid- and late spring bloomers *Iris pumila* L., *Convallaria majalis* L., and *Polygonatum multiflorum* L. To maximize genetic conformity, vegetatively propagated specimens from the same location were used. The specimens were grown in standardized pots with a diameter of 14 cm, and were signed individually. All selected species were pollinated by insects. 5 repetitions were used for each species at both sites.

The *ex situ* experiment were set-up between December 2019 and February 2020. The *Galanthus nivalis* and *Eranthis hyemalis* specimens were introduced in autumn 2020, so we have data from 2021. The same soil mixture and irrigation protocol was used for every specimen at both sites. The plants were watered twice a week in spring and autumn and daily in summer to keep them well hydrated.

Measurements were taken for each specimens on the same day and weekly basis at both sites for 3 consecutive years. The experiment period covers the interval between March of 2020 and December of 2022.

We used the average temperature difference (1.81 °C) between the two locations, measured during the three years of the experiment, to calculate the phenological sensitivity. It means the difference in the appearance of a phenophase (the time interval) was divided by 1.81 to get the phenological sensitivity. For example in case of one day difference the phenological sensitivity is  $1 \text{ day} / 1.81 \text{ °C} = 0.55 \text{ day/°C}$ .

## 2.3. Measurements of biotic data

The species used for the experiment all have solitary flower. We recorded the number of buds, flowers and fruits by every uniquely signed plant specimen (numbered pots), at both site on a weekly basis.

## 2.4 Measurements of abiotic data (environmental parameters)

The meteorological data were collected by AgroSense base weather station (Sys-Control Kft, Budapest, Hungary) installed at both sites at the end of 2020 (*Figure 1*). For the year 2020 the data (daily mean temperature and daily sum of precipitation) of the nearest station of the Hungarian Meteorological Service (Lágymányos for Budapest and Aszód for Gödöllő) were used.




Figure 1: Climatic conditions of the sites during the duration of the experiment (March 2020 – Dec. 2022): monthly variation of mean air temperature (a) and sum of precipitation (b)


### 2.5. Statistical analysis

Data recording and basic data compilation was carried out in Microsoft Excel 365 online version and all statistical analyses were performed using freely available software R, version 4.2.2. (R Core Team R 2022), together with RStudio script editor (RStudio Team 2015). For advanced data processing, the additional packages “tidyverse” (Wickham et al. 2019), “dplyr” (Wickham et al. 2023) and “scales”

(Wickham et al. 2022) were also used. Package “ggplot2” (Wickham et al. 2016) was used for creating advanced statistical graphs.


### 3. Results

The relatively short flowering period of the species, and occasional non-flowering or mortality, resulted in relatively low element counts.



*Figure 2:* Date of beginning of flowering (first flower) at both sites. The letters **a** and **b** are for the mean values (green numbers), differing letters indicate a statistical significance difference at 5% significance level. The letter **n** is for the sample size. Standard deviation is shown on the top of the columns.

The appearance of the first flower (*Figure 2*) occurred earlier for 4 of the 5 species in all three years at the warmer Budapest site. The difference is significant in 2 species. One species, *Eranthis hyemalis*, showed the opposite effect, the beginning of flowering was earlier at the Gödöllő site, but the difference is not significant. In the case of this species, the result is influenced by the fact that the individuals of the species died at the Budapest site by 2022, so they did not bloom here that year at all, while they did in Gödöllő. On average, the time of the appearance of the first flower at the Budapest location is day of year (DOY) 95.20, while at the Gödöllő location DOY 98.84, the difference is 3.64 days. A virágzás hossza átlagosan 10.52 nap volt a budapesti helyszínen, míg 8.22 a gödöllői helyszínen, the difference is 2.3 days. Phenological sensitivity: 2.01 day/°C.



*Figure 3: End of flowering.* The letters **a** and **b** are for the mean values (green numbers), differing letters indicate a statistical significance difference at 5% significance level. The letter **n** is for the sample size. Standard deviation is shown on the top of the columns.

The end of flowering (*Figure 3*) occurred earlier for 3 of the 5 species in all three years at the warmer Budapest site. The difference is significant in 2 species – with earlier end of flowering at the Budapest site, and in one species with earlier end of flowering at the Gödöllő site.

On average, the time of the end of the flowering at the Budapest location is DOY 105.72 while at the Gödöllő location DOY 107.06, the difference is 1.34 days. Phenological sensitivity: 0.74 day/°C. On average, the length of flowering at the Budapest location was 10.52 days, while at the Gödöllő location 8.22 days, the difference is 2.3 days. Phenological sensitivity: 1.27 day/°C.

#### 4. Discussion

Considering all species, the first flower appeared at the Budapest location on DOY 95.20, while at the Gödöllő location on DOY 98.84. The difference is 3.64 days which indicates a phenological sensitivity of 2.01 days/°C. On average, the end of the flowering occurred at the Budapest site on DOY 105.72 while at the Gödöllő site on DOY 107.06. The difference is 1.34 days, which indicates a phenological sensitivity of 0.74 days/°C. The difference between the sensitivity of the beginning and the end of flowering amounts to a difference of 2.72 times, where the stronger one is the sensitivity of the beginning of the flowering. It means, that the length of flowering was in Budapest 10,52 days long, while in Gödöllő 8,22 days long. The difference between the phenological sensitivity of the beginning and the end of

flowering results in a longer flowering period at the site with a higher average temperature. This is in contrast to previous research (Miller-Rushing et al. 2009, Sherry et al. 2011, Bock et al. 2014, Nagahama et al. 2018), which found that higher temperatures shorten flowering time.

However, it is worth examining the early bloomer species (*Galanthus nivalis* and *Eranthis hyemalis*) and the mid- and late-bloomer species (*Iris pumila*, *Convallaria majalis* and *Polygonatum multiflorum*) separately. Regarding the early bloomer geophytes, the first flower appeared on average on DOY 62.75 in Gödöllő and on DOY 64.9 in Budapest. It means, the flowering advanced 2.15 days at the site with the lower average temperature, in contrary to previous researches (Sparks et al. 2000, Fitter and Fitter 2002, Parmesan and Yohe 2003, Elzinga et al. 2007, Bertin 2008, Szabó et al. 2016, Neumann & Czóbel 2021). There are several reasons for this. On the one hand, there was little data available, as the early geophytes only arrived in the second year of the experiment, and on the other hand, the fact that the early geophyte specimens at the Budapest site were dead by 2022 may have put a damper on the data. If we consider only the year 2021, when both early species flowered at both sites, the onset of flowering is DOY 54 and 47.5 in Budapest, values that are consistent with the general trend of earlier flowering in areas with higher average temperatures (Sparks et al. 2000, Fitter and Fitter 2002, Parmesan and Yohe 2003, Elzinga et al. 2007, Bertin 2008, Szabó et al. 2016, Neumann & Czóbel 2021).

Fitter et al. (1995) found that early-flowering species are very variable, regarding the flowering onset dates. The early bloomers showed the end of flowering on DOY 79.35 in Gödöllő and on 73.5 in Budapest. It results in a length of flowering of 16.6 days in Gödöllő and 8.6 days in Budapest. It means that the flowering period was shorter at the site with the higher average temperature which is in line with the previous research (Miller-Rushing et al. 2009, Sherry et al. 2011, Bock et al. 2014, Nagahama et al. 2018). It means a phenological sensitivity of 4,42 days/°C.

Regarding the mid- and late bloomer geophytes, the first flower appeared on average on DOY 122.9 in Gödöllő and on DOY 115.4 in Budapest. The difference is 7.5 days, which indicates a phenological sensitivity of 4.14 days/°C. It is in line with previous research, Fitter et al. (1995) found that high spring temperatures advanced flowering by a mean of 4 days per degree. The end of flowering occurred on DOY 125.5 in Gödöllő and on DOY 127.2 in Budapest. The difference is 1,7 days, surprisingly with earlier occurrence at the Gödöllő site with the lower average temperature. This indicates a phenological sensitivity of -0,94 days/°C. The length of flowering was in Budapest 11,8 days long, while in Gödöllő 2,6 days long. This is in contrast to previous research (Miller-Rushing et al. 2009, Sherry et al. 2011, Bock et al. 2014, Nagahama et al. 2018), which found that higher temperatures shorten flowering time.

Previous research found that ephemerals, early-spring bloomers, and insect-pollinated plants in these environments tend to be more sensitive and show a greater advancement than perennials, mid- or late-spring bloomers, and wind-pollinated plants. It is in line with other researches (Fitter and Fitter 2002, Nail and Wu 2006). To find out, a longer-term, multi-species study would be needed.



Changes in the phenology of different native geophyte species due to climate change will bring about various other changes and will challenge native conservation. De Frenne et al. (2011) found that understorey species will probably advance with increasing temperatures in the future, the effects on growth and reproductive performance are species-dependent. It is in line with our findings. de Frenne et al. (2011) concluded that these divergent responses of understorey plants could alter future forest understorey dynamics included community assembly. The phenological shift can cause several problems, some also related to the nature conservation. Zettlemoyer et al. (2019) found that warming-led non-native species were likely to flower earlier and more plastic to temperature than the natives. Phenology will alter temporal overlap between plants and pollinators. can cause mismatches between plants and their pollinators (Forrest 2015). Warming caused shortened winter periods, alongside decreasing snow cover duration, increase late spring frosts frequency (Liu et al., 2018; Zohner et al., 2020), which can negatively affect plant growing conditions in spring.

### Acknowledgments

The research was supported by the Doctoral School of Environmental Sciences of the Hungarian University of Agriculture and Life Sciences. We are grateful Ildikó Pándy, the Hungarian University of Agriculture and Life Sciences' head of Gödöllő Botanical Garden, and László Papp, botanist in ELTE Botanical Garden, for the professional and technical help during experimental set-up. Special thank goes to Enikő Szentpéteri and Márk Pozsonyi for the enormous help by taking care of the plants during the experiment.

### References

- Bandoc G., Piticar A., Patriche C., Roșca B., Dragomir E.(2022): Climate Warming-Induced Changes in Plant Phenology in the Most Important Agricultural Region of Romania. Sustainability. 14: 2776. <https://doi.org/10.3390/su14052776>
- Bertin Robert I. (2008): "Plant Phenology and Distribution in Relation to Recent Climate Change." The Journal of the Torrey Botanical Society. 135 (1): 126–46. JSTOR, <http://www.jstor.org/stable/20063966>.
- Blois J.L., Williams J.W., Fitzpatrick M.C., Jackson S.T., Ferrier S. (2013): Space can substitute for time in predicting climate-change effects on biodiversity. Proc. Natl. Acad. Sci. 110, 9374–9379. 1220228110 <https://doi.org/10.1073/pnas.1220228110>
- Bock A., Sparks T. H., Estrella N., Jee N., Casebow A., Schunk C., Menzel A. (2014): Changes in first flowering dates and flowering duration of 232 plant species on the island of Guernsey. Global Change Biology. 20(11): 3508–3519. doi:10.1111/gcb.12579
- Cleland EE., Chuine I., Menzel A., Mooney HA., Schwartz MD.(2007): Shifting plant phenology in response to global change. Trends Ecol Evol. 22 (7):357-65. doi: 10.1016/j.tree.2007.04.003.
- Crișan I., Stoie A., Buta E., Cantor M.(2018): Flowering phenology of some Iris species in the UASVM Cluj agrobotanical garden, Romanian Biotechnological Letters, 2018, Vol. 23, No. 3, 2018
- De Frenne P., Brunet J., Shevtsova A., Kolb A., J Graae B., Chabrerie O., Ao Cousins S., Decocq G., De Schrijver A., Diekmann M., Gruwez R., Heinken T., Hermy M., Nilsson C., Stanton S., Tack W., Willaert J., Verheyen K. (2011): Temperature effects on forest herbs assessed by

## Review on Agriculture and Rural Development 2023 vol. 12 (1-2)

- warming and transplant experiments along a latitudinal gradient, *Global Change Biology*, vol. 17, issue 10, pp. 3240-3253, 2011, doi: 10.1111/j.1365-2486.2011.02449.x
- Elzinga J.A., Atlan A., Biere A., Gigord L., Weis A.E., Bernasconi G. (2007): Time after time: flowering phenology and biotic interactions. *Trends Ecol Evol.* 22 (8): 432-9. doi: 10.1016/j.tree.2007.05.006. Epub 2007 Jun 15. PMID: 17573151.
- Eppich B., Dede L., Ferenczy A., Garamvölgyi Á., Horváth L., Isépy I., Priszter Sz, Hufnagel L. (2009): Climatic effects on the phenology of geophytes. *Applied Ecology and Environmental Research.* 7. 253-266. [https://doi.org/10.15666/aeer/0703\\_253266](https://doi.org/10.15666/aeer/0703_253266)
- Fitter AH., Fitter RS.(2002): Rapid changes in flowering time in British plants. *Science.* 31: 296(5573):1689-91. doi: 10.1126/science.1071617. PMID: 12040195.
- Fitter A.H., Fitter R. S. R., Harris I. T. B., Williamson M. H. (1995): Relationships Between First Flowering Date and Temperature in the Flora of a Locality in Central England. *Functional Ecology.* 9 (1): 55–60. JSTOR, <https://doi.org/10.2307/2390090>.
- Forrest J., Miller-Rushing AJ.(2010): Toward a synthetic understanding of the role of phenology in ecology and evolution. *Philos Trans R Soc Lond B Biol Sci.* 2010 Oct 12;365(1555):3101-12. doi: 10.1098/rstb.2010.0145. PMID: 20819806; PMCID: PMC2981948.
- Forrest J.R.K. (2015): Plant–pollinator interactions and phenological change: what can we learn about climate impacts from experiments and observations?. *Oikos*, 2015, 124: 4-13. <https://doi.org/10.1111/oik.01386>
- IPCC 2022: *Climate Change 2022: Impacts, Adaptation and Vulnerability. Contribution of Working Group II to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* [H.-O. Pörtner, D.C. Roberts, M. Tignor, E.S. Poloczanska, K. Mintenbeck, A. Alegría, M. Craig, S. Langsdorf, S. Lösschke, V. Möller, A. Okem, B. Rama (eds.)]. Cambridge University Press. Cambridge University Press, Cambridge, UK and New York, NY, USA, 3056 pp., doi:10.1017/9781009325844.
- Khodorova NV., Boitel-Conti M. (2013): The Role of Temperature in the Growth and Flowering of Geophytes. *Plants (Basel).* 2(4):699-711. doi: 10.3390/plants2040699. PMID: 27137399; PMCID: PMC4844387.
- Kubov M., Schieber B. Janík R.(2022): Effect of Selected Meteorological Variables on Full Flowering of Some Forest Herbs in the Western Carpathians. *Atmosphere* 195. <https://doi.org/10.3390/atmos13020195>
- Liu Q., Piao S., Janssens I.A., Fu Y., Peng S., Lian X., Ciais P., Myneni R.B., Peñuelas J., Wang T.: (2018): Extension of the growing season increases vegetation exposure to frost. *Nat. Commun.*, 9: 426. <https://doi.org/10.1038/s41467-017-02690-y>.
- Menzel A., Sparks T. H., Estrella N., Koch E., Aasa A., Ahas R., Almkübler K., Bissolli P., Braslavská O., Briede A., Chmielewski F. M., Crepinsek Z., Curnel Y., Dahl Å., Defila C., Donnelly A., Filella Y., Jatczak K., Måge F., Mestre A., Nordli Ø., Peñuelas J., Pirinen P., Remišová V., Scheifinger H., Striz M., Susnik A., van Vliet AJH., Wielgolaski FE., Zach S., Züst A. (2006): European phenological response to climate change matches the warming pattern. *Global Change Biology*, 12, 1969–1976. <https://doi.org/10.1111/j.1365-2486.2006.01193.x>
- Miller-Rushing A. J., & Primack R. B. (2008): Global warming and flowering times in Thoreau's concord: A community perspective. *Ecology*, 89: 332–341. <https://doi.org/10.1890/07-0068.1>
- Miller-Rushing A.J. & Inouye D.W. (2009): Variation in the impact of climate change on flowering phenology and abundance: An examination of two pairs of closely related wildflower species. *American Journal of Botany*, 96: 1821-1829. <https://doi.org/10.3732/ajb.0800411>
- Nagahama A., Kubota Y. & Satake A. (2018): Climate warming shortens flowering duration: a comprehensive assessment of plant phenological responses based on gene expression analyses and mathematical modeling. *Ecological Research*, 33: 1059–1068. <https://doi.org/10.1007/s11284-018-1625-x>
- Neil K., Wu J. (2006): Effects of urbanization on plant flowering phenology: A review. *Urban Ecosystem*, 9: 243–257. <https://doi.org/10.1007/s11252-006-9354-2>

## Review on Agriculture and Rural Development 2023 vol. 12 (1-2)

- Orlóci L.; Kiszél P.; Solymosiné László I.; Papp L. (2019): *Delectus Seminum Sporarum Plantarumque Horti Botanici Universitatis Hungariae*. Eötvös Lóránd Tudományegyetem. Botanikus Kertje Universitatis Scientiarum Hungariae de Lorand Eoetvoes Nuncupatae, 165th ed.; Fűvészkeret Alapítvány (Foundation): Budapest, Hungary, 36p.
- Peñuelas J., Ciais P., Canadell J.G. et al. (2017): Shifting from a fertilization-dominated to a warming-dominated period. *Nature Ecology and Evolution*, 1: 1438–1445, <https://doi.org/10.1038/s41559-017-0274-8>
- Pickett S.T.A. (1989): Space-for-Time Substitution as an Alternative to Long-Term Studies. In: Likens G.E. (eds) *Long-Term Studies in Ecology*. Springer, New York, NY. [https://doi.org/10.1007/978-1-4615-7358-6\\_5](https://doi.org/10.1007/978-1-4615-7358-6_5)
- R Core Team R. *A Language and Environment for Statistical Computing*; R Foundation for Statistical Computing: Vienna, Austria, (2022); Available online: <http://www.R-project.org/> (accessed on 15 March 2023).
- Puchałka R., Klisz M., Koniakin S., Czortek P., Dylewski Ł., Paż-Dyderska S., Vítková M., Sádlo J., Rašomavičius V., Čarni A., De Sanctis M., Dyderski M. K., (2022): Citizen science helps predictions of climate change impact on flowering phenology: A study on *Anemone nemorosa*, *Agricultural and Forest Meteorology*, 325: (2022) 109133, ISSN 0168-1923, <https://doi.org/10.1016/j.agrformet.2022.109133>.
- Renner S. S., & Zohner C. M. (2018). Climate change and phenological mismatch in trophic interactions among plants, insects, and vertebrates. *Annual Review of Ecology, Evolution, and Systematics*, 49, 165–182. <https://doi.org/10.1146/annurev-ev-ecolsys-110617-062535>
- RStudio Team. *RStudio: Integrated Development for R*; RStudio Inc.: Boston, FL, USA, 2015; Available online: <http://www.rstudio.com/> (accessed on 15 March 2023).
- Sherry Rebecca A. et al. (2007) Divergence of reproductive phenology under climate warming, *J Proceedings of the National Academy of Sciences*, P 198-202, V 104, N 1, doi:10.1073/pnas.0605642104
- Sherry R. A., Zhou X., Gu S., Arnone J. A. III, Johnson D. W., Schimel D. S., Verburg P. S.J., Wallace L. L. & Luo Y. (2011): Changes in duration of reproductive phases and lagged phenological response to experimental climate warming, *Plant Ecology & Diversity*, 4:(1) 23-35, DOI: 10.1080/17550874.2011.557669
- Snopková Z., Hýrošová T. (2017): Snow cover and its influence on the beginning of flowering of snowdrop (*Galanthus nivalis* L.) at the international phenological station (gpm) in Banská Bystrica over the period from 2003 to 2017 In *Snow an ecological phenomenon*, Smolenice, Slovakia, 19th – 21st September 2017.
- Sparks T., Jeffree E. & Jeffree C. (2000): An examination of the relationship between flowering times and temperature at the national scale using long-term phenological records from the UK. *Int J Biometeorol* 44: 82–87. <https://doi.org/10.1007/s004840000049>
- Sparks T.H. and Menzel A. (2002): Observed changes in seasons: an overview. *Int. J. Climatol.*, 22: 1715-1725. <https://doi.org/10.1002/joc.821>
- Stewart D., Oke T.R. (2012) *Local Climate Zones for Urban Temperature Studies*: Bulletin of the American Meteorological Society Volume 93, Issue 12: 1879–1900 <https://doi.org/10.1175/BAMS-D-11-00019.1>
- Szabó B., Vincze E., Czúcz B. (2016): Flowering phenological changes in relation to climate change in Hungary. *Int J Biometeorol*. 60(9):1347-56. doi: 10.1007/s00484-015-1128-1. Epub 2016 Jan 14. PMID: 26768142.
- Szirmai O., Horel J., Neményi A., Pándi I., Gyuricza C.S., Czóbel Sz. (2014): Overview of the collections of the first agrobotanical garden of Hungary. *Hung. Agric. Res.* 23: 19–25.
- Tarasjev A. (1997), Flowering phenology in natural populations of *Iris pumila*. *Ecography*, 20: 48–54. <https://doi.org/10.1111/j.1600-0587.1997.tb00346.x>
- Thomson JD. (2010): Flowering phenology, fruiting success and progressive deterioration of pollination in an early-flowering geophyte. *Philos Trans R Soc Lond B Biol Sci.* 2010 Oct 12;365(1555):3187-99. doi: 10.1098/rstb.2010.0115. PMID: 20819812; PMCID: PMC2981941.

## Review on Agriculture and Rural Development 2023 vol. 12 (1-2)

- Turisova I., Snopková Z., Škvareninová J. (2007). Priestorová Analýza Nástupu Začiatku Kvitnutia Convallaria Majalis L. Na Strednom Slovensku. 114. 978-80.
- Verbényiné Neumann, K.; Czóbel Sz. (2021): Comparative study of flowering phenology of selected plant life forms in urban and rural environments. Preliminary results, pp. 25-36 Columella — Journal of Agricultural and Environmental Sciences Vol. 8. No.1 (2021) p. 65, DOI: 10.18380/SZIE.COLUM.2021.8.1.25
- Wickham H. Ggplot2: Elegant Graphics for Data Analysis; Springer: New York, NY, USA, 2016; 213p.
- Wickham H., Averick M., Bryan J., ChangW., McGowan L., François R., Grolemond G., Hayes A., Henry L., Hester J. et al. (2019): Welcome to the tidyverse. J. Open Source Softw, 4: 1686.
- Wickham H., Francois R., Henry L., Müller K., Vaughan D. Dplyr (2023): A Grammar of Data Manipulation. R Package Version 1.1.0. 2023. Available online: <https://CRAN.R-project.org/package=dplyr> (accessed on 15 March 2023).
- Wickham H., Seidel D. Scales (2022): Scale Functions for Visualization. R Package Version 1.2.1. 2022. Available online: <http://CRAN.R-project.org/package=scales> (accessed on 15 March 2023).
- Wolkovich E., Donahue MJ. (2021): How phenological tracking shapes species and communities in nonstationary environments., Biol. Rev., 96: 2810–2827. doi:10.1111/brv.12781
- Zettlemoyer MA., Schultheis EH., Lau JA. (2019): Phenology in a warming world: differences between native and non-native plant species. Ecol Lett. 2019, xxx, <https://doi.org/10.1111/ele.13290>
- Ziska LH., Gebhard DE., Frenz DA., Faulkner S., Singer BD., Straka J.: (2003) Cities as harbingers of climate change: Common ragweed, urbanization, and public health. J Allergy Clini Immun 111:290–295 <https://doi.org/10.1067/mai.2003.53>
- Zohner C.M., Mo, L., Renner S.S., Svenning J.C., Vitasse Y., Benito B.M., Ordonez A., Baumgarten F., Bastin J.F., Sebald V., Reich P.B., Liang J., Nabuurs G.J., De- Migueln S., Alberti G., Ant´on-Fern´andez C., Balazy R., Br´andli U.B., Chen H.Y.H., Chisholm C., Cienciala E., Dayanandan S., Fayle T.M., Frizzera L., Gianelle D., Jagodzinski A.M., Jaroszewicz B., Jucker T., Kepfer-Rojas S., Khan M.L., Kim H. S., Korjus H., Johannsen V.K., Laarmann D., Langn M., Zawila-Niedzwiecki T., Niklaus P.A., Paquette A., Pretzsch H., Saikia P., Schall P., Seben V., Svoboda M., Tikhonova E., Viana H., Zhang C., Zhao X., Crowther T.W. (2020): Late-spring frost risk between 1959 and 2017 decreased in North America but increased in Europe and Asia. Proc. Natl. Acad. Sci. U. S. A. 117, 12192–12200. <https://doi.org/10.1073/pnas.1920816117>.

Electronic reference:

[http1:https://geopedia.world/#T4\\_L107\\_x2124674.6907575\\_y6021482.165878462\\_s16\\_b2345](http1:https://geopedia.world/#T4_L107_x2124674.6907575_y6021482.165878462_s16_b2345)  
(downloaded 26 April 2021)